Measurement of Inclusive Jet Cross Sections in $Z/\gamma^*(\rightarrow e^+e^-)$+jets Production in $p\overline{p}$ Collisions at $\sqrt{s}$ = 1.96 TeV

M. D'Onofrio, M. Martínez, O. Saltó (for the CDF collaboration)
IFAE/ICREA, Institut de Física d'Altes Energies, Barcelona, E-08193, Spain.

Inclusive jet cross sections in Z/γ^* events, with Z/γ^* decaying into an electron-positron pair, are measured as a function of jet transverse momentum and jet multiplicity in $p\overline{p}$ collisions at $\sqrt{s} = 1.96$ TeV with the upgraded Collider Detector at Fermilab in Run II, based on an integrated luminosity of 2.5 fb^{-1}. The measurements cover the rapidity region $|y^{\rm jet}| < 2.1$ and the transverse momentum range $p_{\rm T}^{\rm jet} > 30$ GeV/c. Next-to-leading order perturbative QCD predictions are in good agreement with the measured cross sections.

arXiv:0810.3642v2 [hep-ex] 20 Oct 2008

1. INTRODUCTION

The study of the production of electroweak bosons in association with jets of hadrons in the final state constitutes a fundamental item in the high-$p_{\rm T}$ physics program at the Tevatron. These events are main backgrounds to many interesting physics processes like, for example, top production, the search for the SM Higgs, and supersymmetry. The CDF experiment has recently published precise measurements on jets in events with a Z/γ^* boson in the final state [1], where inclusive jet cross sections as a function of jet transverse momentum and jet multiplicity are measured and compared to pQCD predictions [2]. At the leading order (LO) in pQCD, Z/γ^*+jet events are driven by the processes $gq \rightarrow Z/\gamma^* + q$ and $q\overline{q} \rightarrow Z/\gamma^* + g$, while higher orders contributions, including additional parton radiation, produce mutiple jets in the final state. Next-to-leading order (NLO) pQCD predictions for Z/γ^*+jets production are only available for jet multiplicities $N_{\rm jet}$ up to $N_{\rm jet} = 2$. This contribution presents updated results with an increased data sample corresponding to a total integrated luminosity of 2.5 fb^{-1}.

2. EVENT SELECTION

The events are required to have two electrons with $E_T^e > 25$ GeV and a reconstructed invariant mass in the range $66 < M_{ee} < 116$ GeV/c^2 around the Z boson mass. The electron candidates are reconstructed using criteria described in [3]. In this study, one electron is required to be central ($|\eta^e| < 1$) and fulfill tight selection cuts, while the second electron is required to pass a looser selection and to be either central or forward with $1.2 < |\eta^e| < 2.8$. The events are selected to have a reconstructed primary vertex with z-position within 60 cm around the nominal interaction point, and at least one jet with transverse momentum $p_{\rm T}^{\rm jet} > 30$ GeV/c, rapidity in the range $|y_{\rm cal}^{\rm jet}| < 2.1$, and $\Delta R_{e-jet} > 0.7$, where ΔR_{e-jet} denotes the distance between the jet and each of the two electrons in the final state. The main backgrounds to the $Z/\gamma^*(\rightarrow e^+e^-)$+jets sample arise from inclusive-jets and W+jets events, and are estimated from the data. Other background contributions from $t\overline{t}$, $Z/\gamma^*(\rightarrow e^+e^-) + \gamma$, WW, WZ, ZZ, and $Z/\gamma^*(\rightarrow \tau^+\tau^-)$+jets final states are estimated using Monte Carlo samples. The total background in inclusive $Z/\gamma^*(\rightarrow e^+e^-)$+jets production is about 12% for $N_{\rm jet} \geq 1$, and increases up to about 17% for $N_{\rm jet} \geq 3$.

3. UNFOLDING PROCEDURE

The measured cross sections are corrected for acceptance and smearing effects back to the hadron level using PYTHIA-TUNE A Monte Carlo event samples [4, 5], CTEQ5L [6] parton distribution functions (PDFs) for the proton and antiproton, and a bin-by-bin unfolding procedure that also accounts for the efficiency of the $Z/\gamma^*(\rightarrow e^+e^-)$ selection criteria. The final results refer to hadron level jets with $p_{\rm T}^{\rm jet} > 30$ GeV/c and $|y^{\rm jet}| < 2.1$, in a limited and well-defined kinematic range for the Z/γ^* decay products: $E_T^e > 25$ GeV, $|\eta^{e1}| < 1.0$, $|\eta^{e2}| < 1.0$ or $1.2 < |\eta^{e2}| < 2.8$,



Figure 1: (top)Measured inclusive jet differential cross section as a function of $p_{\mathrm{T}}^{\mathrm{jet}}$ (black dots) in $Z/\gamma^*(\rightarrow e^+e^-)$+jets with $N_{\mathrm{jet}} \geq 1, 2$ compared to NLO pQCD predictions (open circles). For clarity, the measurement for $N_{\mathrm{jet}} \geq 1$ is scaled up by ($\times$20). The shaded bands show the total systematic uncertainty, except for the 5.8% luminosity uncertainty. (middle and bottom) Data/theory ratio as a function of $p_{\mathrm{T}}^{\mathrm{jet}}$ for $N_{\mathrm{jet}} \geq 1$ and $N_{\mathrm{jet}} \geq 2$, respectively. The dashed and dotted lines indicate the PDF uncertainty and the variation with μ of the NLO pQCD predictions, respectively.

$66 < M_{ee} < 116$ GeV/c^2, and $\Delta R_{e-jet} > 0.7$. In order to avoid any bias on the correction factors due to the particular PDF set used, which translates into slightly different simulated $p_{\mathrm{T,cal}}^{\mathrm{jet}}$ distributions, the PYTHIA-TUNE A Monte Carlo event sample is re-weighted until it accurately follows the measured $p_{\mathrm{T,cal}}^{\mathrm{jet}}$ spectra. The unfolding factors $U(p_{\mathrm{T,cor}}^{\mathrm{jet}}) = \frac{\mathrm{d}\sigma}{dp_{\mathrm{T}}^{\mathrm{jet}}} / \frac{\mathrm{d}\sigma}{\mathrm{dp}_{\mathrm{T,cor}}^{\mathrm{jet}}}$ are computed separately for the different measurements and vary between 2.0 at low $p_{\mathrm{T}}^{\mathrm{jet}}$ and 2.3 at high $p_{\mathrm{T}}^{\mathrm{jet}}$.

4. COMPARISON WITH PQCD PREDICTIONS

Figure 1(top) shows the measured inclusive jet differential cross sections as a function of $p_{\mathrm{T}}^{\mathrm{jet}}$ in $Z/\gamma^*(\rightarrow e^+e^-)$+jets production, with $N_{\mathrm{jet}} \geq 1$ and $N_{\mathrm{jet}} \geq 2$, compared to NLO pQCD predictions. The cross sections decrease by more than three orders of magnitude as $p_{\mathrm{T}}^{\mathrm{jet}}$ increases from 30 GeV/c up to about 300 GeV/c. The NLO pQCD predictions are computed using the MCFM program [2] with CTEQ6.1M PDFs [7], with the renormalization and factorization scales set to $\mu^2 = M_Z^2 + p_T^2(Z)$, and using a midpoint [8] algorithm with $R = 0.7$ and $R_{\mathrm{sep}} = 1.3$ [9] to reconstruct jets at the parton level. The theoretical predictions include parton-to-hadron correction factors $C_{\mathrm{had}}(N_{\mathrm{jet}}, p_{\mathrm{T}}^{\mathrm{jet}})$ that approximately account for non-perturbative contributions from the underlying event and fragmentation into hadrons. In each measurement C_{had} is estimated using the PYTHIA-TUNE A Monte Carlo samples, as the ratio between the nominal $p_{\mathrm{T}}^{\mathrm{jet}}$ distribution and the one obtained by turning off both the interactions between proton and antiproton remnants and the string fragmentation in the Monte Carlo samples. The correction decreases as $p_{\mathrm{T}}^{\mathrm{jet}}$ increases from about 1.2 (1.26) at $p_{\mathrm{T}}^{\mathrm{jet}}$ of 30 GeV/c to 1.02 (1.01) for $p_{\mathrm{T}}^{\mathrm{jet}} > 200$ GeV/c for $N_{\mathrm{jet}} \geq 1$ ($N_{\mathrm{jet}} \geq 2$), and is dominated by the underlying event contribution. The uncertainty on C_{had} is about 10% (17%) at low $p_{\mathrm{T}}^{\mathrm{jet}}$ and goes down to 1% at high $p_{\mathrm{T}}^{\mathrm{jet}}$ for $N_{\mathrm{jet}} \geq 1$ ($N_{\mathrm{jet}} \geq 2$). The ratios between data and theory as a function of $p_{\mathrm{T}}^{\mathrm{jet}}$ are also shown in Fig. 1.

Good agreement is observed between the measured cross sections and the nominal theoretical predictions.



Figure 2: (top) Measured cross section for inclusive jet production in $Z/\gamma^*(\rightarrow e^+e^-)$ events as a function of $N_{\rm jet}$ compared to LO and NLO pQCD predictions. The shaded bands show the total systematic uncertainty, except for the 5.8% luminosity uncertainty. (bottom) Ratio of data and NLO to LO pQCD predictions versus $N_{\rm jet}$. The dashed and dotted lines indicate the PDF uncertainty and the variation with μ of the NLO pQCD predictions, respectively.

Finally, Fig. 2 shows the cross sections $\sigma_{N_{\rm jet}}$ for $Z/\gamma^*(\rightarrow e^+e^-)$+jets events up to $N_{\rm jet} \geq 3$. The data are compared to LO and NLO pQCD predictions. The parton-to-hadron non-perturbative corrections vary between 1.1 and 1.4 as $N_{\rm jet}$ increases. The LO pQCD predictions underestimate the measured cross sections by a factor about 1.4 approximately independent of $N_{\rm jet}$. Good agreement is observed between data and NLO pQCD predictions.

The final data sample collected by the CDF experiment in Run II, where more than 6 fb^{-1} are expected, will make possible further detailed studies of the event topologies.

Acknowledgments

We would like to thank the organizers for their kind invitation to the conference, the wonderful atmosphere, and the discussions that they made possible.

References

[1] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **100**, 102001 (2008).
[2] J. Campbell and R.K. Ellis, Phys. Rev. D **65**, 113007 (2002).
[3] A. Abulencia *et al.* (CDF Collaboration), J. Phys. G: Nucl. Part. Phys. **34**, 2457 (2007). Charge conjugation is implied throughout the paper.
[4] T. Sjöstrand *et al.*, Comp. Phys. Comm. **135**, 238 (2001).
[5] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. D **65**, 092002 (2002).
[6] H.L. Lai *et al.*, Eur. Phys. J. C **12**, 375 (2000).
[7] J. Pumplin *et al.*, J. High Energy Phys. **0207**, 012 (2002).
[8] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **74**, 071103(R) (2006).
[9] S.D. Ellis and D.E. Soper, Phys. Rev. D **48**, 3160 (1993).